

Dipenguin-like contributions to $D^0 - \bar{D}^0$ mixing

Alexey A. Petrov

*Department of Physics and Astronomy
The Johns Hopkins University
Baltimore MD 21218*

Abstract. We show that because of the milder GIM suppression, the four-quark dipenguin operator contributes to the short distance piece of the $D^0 - \bar{D}^0$ mixing amplitude at the same order of magnitude as the box diagram. In addition, we put an upper bound $\Delta m_D^{LDR} \leq 5 \times \Delta m_D^{box}$ on a long-distance resonant contribution induced by the penguin mixing.

The phenomenon of meson mixing has been studied for a long time. Observed in the $K^0 - \bar{K}^0$ and $B^0 - \bar{B}^0$ systems, it provides an extremely sensitive test of the Standard Model (SM). Extremely large value of the top quark mass makes the short distance box diagram (e.g. Fig.1a) dominate $B^0 - \bar{B}^0$ and $K^0 - \bar{K}^0$ mixing amplitudes. This fact effectively turns GIM cancellation mechanism (vanishing of the total $\Delta Q = 2$ contribution in the limit of equal internal quark masses) into the "GIM-enhancement" of the short-distance contribution to Δm_P for $P = \{K, B\}$. Long-distance contributions associated with the propagation of the hadronic degrees of freedom and higher-order QCD effects are relatively unimportant for the mixing of B and K mesons.

The situation is drastically different in the case of the $D^0 - \bar{D}^0$ system: the only available heavy b -quark contribution to the box diagram providing $\Delta C = 2$ transition is greatly reduced by a tiny V_{ub} CKM matrix element and can be safely neglected in the analysis of the box diagram contribution to Δm_D (the inclusion of the b -quark further decreases the box diagram contribution [1]). This forces the vanishing of the effect in the $SU(3)$ limit and implies that only the light quark mass difference guarantees that mixing does take place. Calculating a box diagram and constructing the effective Hamiltonian, we realize that the smallness of the short distance piece is guaranteed by a factor of $(m_s^2 - m_d^2)/m_c^2$ [2] (we define $L = (1 + \gamma_5)/2$)

$$\mathcal{H}_{eff} = \frac{G_F^2}{2\pi^2} \xi_s \xi_d \frac{(m_s^2 - m_d^2)^2}{m_c^2} [\bar{u} \gamma_\mu L c \bar{u} \gamma_\mu L c + 2 \bar{u} R c \bar{u} R c] \quad (1)$$

with $\xi_i = V_{ic}^* V_{iu}$. We denote the operators, entering Eq. (1) as O_1 and O_2 respectively. This leads to the following expression for the Δm_D^{box} [1,2]

$$\Delta m_D^{box} = \frac{4G_F^2}{3\pi^2} \xi_s \xi_d \frac{(m_s^2 - m_d^2)^2}{m_c^2} f_D^2 m_D (B_D - 2B'_D) \approx 0.5 \cdot 10^{-17} \text{ GeV} \quad (2)$$

with $f_\pi \simeq 132 \text{ MeV}$, $f_D \simeq 165 \text{ MeV}$, $m_s = 0.2 \text{ GeV}$, and $B_D = B'_D = 1$ in the usual vacuum saturation approximation to

$$\langle D^0 | O_1 | \bar{D}^0 \rangle = \frac{2}{3} \frac{f_D^2 m_D^2}{2m_D} B_D, \quad \langle D^0 | O_2 | \bar{D}^0 \rangle = -\frac{5}{3} \frac{m_D^2}{4m_c^2} \frac{f_D^2 m_D^2}{2m_D} B'_D. \quad (3)$$

The smallness of the result of Eq.(2) triggered a set of estimates of the long-distance effects and New Physics contributions that can dominate the Δm_D .

The box diagram, however, does not exhaust the set of short-distance SM contributions. Higher order QCD corrections might significantly modify the leading order prediction provided that they are not proportional to the small factor $(m_s^2 - m_d^2)^2/m_c^2$. For instance, a set of diagrams (topologically distinct from the box diagram) generated by the so-called “dipenguin” operators (Fig.1b.) contribute to the D -meson mass difference at the same order of magnitude as the usual box diagram [3]. To see this, let us write an effective operator relevant to dipenguin $\Delta C = 2$ transition. It can be obtained from the usual $\Delta C = 1$ penguin vertex. Neglecting a tiny dipole contribution,

$$\Gamma = -\frac{G_F}{\sqrt{2}} \frac{g_s}{4\pi^2} F_1 \bar{u} \gamma_\mu L t^a c (g^{\mu\nu} \partial^2 - \partial^\mu \partial^\nu) A_\nu^a. \quad (4)$$

Here $t^a = \lambda^a/2$, and F_1 is a modified Inami-Lim (IL) function. Using unitarity of the CKM matrix it reads $F_1 = \sum_i \xi_i F_1^i = \xi_s (F_1^s - F_1^d) + \xi_b (F_1^b - F_1^d)$. It is common to discard the b -quark contribution to F_1 as being suppressed by small V_{ub} factors. Note, however, that it *enhances* F_1 by approximately 20–30%. From (4) we obtain the effective Hamiltonian

$$\mathcal{H}_{dp} = -\frac{G_F^2}{8\pi^2} \frac{\alpha_s}{\pi} F_1^2 \left[(\bar{u} \gamma_\mu L t^a c) \partial^\mu \partial^\nu (\bar{u} \gamma_\nu L t^a c) - (\bar{u} \gamma_\mu L t^a c) \square (\bar{u} \gamma_\mu L t^a c) \right] \quad (5)$$

In what follows, we denote two operators entering (5) as $\tilde{O}_1$ and $\tilde{O}_2$. Using the equations of motion we obtain for the first operator in (5)

$$\tilde{O}_1 = \bar{u} \gamma_\mu L t^a c \partial^\mu \partial^\nu \bar{u} \gamma_\nu L t^a c \simeq m_c^2 \bar{u} R t^a c \bar{u} R t^a c \quad (6)$$

Employing vacuum saturation method to estimate matrix elements and assuming that each derivative acting on the quark field involves an average momentum of the quark we obtain

$$\langle D^0 | \tilde{O}_1 | \bar{D}^0 \rangle = \frac{1}{9} \frac{f_D^2 m_D^4}{2m_D} B, \quad \langle D^0 | \tilde{O}_2 | \bar{D}^0 \rangle = -\frac{2}{9} \frac{f_D^2 m_D^2 (2m_c^2 - m_D^2)}{2m_D} B' \quad (7)$$




FIGURE 1. Examples of the box diagram (a), dipenguin diagram (b) and a possible long-distance one-particle intermediate state contribution (c).

with B and B' being the bag parameters. This yields (setting $B = B' = 1$)

$$\Delta m_D^{dp} = 2 \langle D^0 | \mathcal{H}^{dp} | \bar{D}^0 \rangle = \frac{G_F^2}{72\pi^2} \frac{\alpha_s}{\pi} F_1^2(m_b^2, m_s^2, m_d^2) f_D^2 m_D (m_D^2 - 4m_c^2). \quad (8)$$

Inspecting (8) we notice that contrary to Δm_D^{box} , the dipenguin diagram does not have a power dependence upon the internal quark masses. The leading behavior of the IL function is logarithmic in $m_{s(d)}$, and the estimate of the operator brings about power dependence upon the *external* quark masses, i.e. m_c . This feature distinguishes this contribution from that of the usual box diagram. Recalling the fact that the dominant contribution to K and B mixing amplitudes is proportional to the square of the top quark mass, it is not surprising that this effect is small in the K and B sectors. It is the fact of the reduced internal heavy quark dependence of the amplitude of $D^0 - \bar{D}^0$ mixing which makes the dipenguin operator contribution effectively enhanced. Taking into account the momentum flow $Q^2 \simeq -m_c^2$ through the gluon propagator (which is defined by the external quark momenta(masses)) one can calculate $F_1^i(m_i^2, Q^2) = -4 \int_0^1 dx x(1-x) \ln[(m_i^2 - Q^2 x(1-x))/M_W^2]$ Comparing (2) and (8) we find [3]

$$r = \left| \frac{\Delta m_D^{dp}}{\Delta m_D^{box}} \right| \approx \frac{\alpha_s}{8\pi} \frac{[F_1(m_b^2, m_s^2, m_d^2)]^2 m_D^4}{|\xi_s \xi_d| (m_s^2 - m_d^2)^2} \quad (9)$$

where we have put $m_c \approx m_D$, and $\alpha_s \simeq 0.4$. The relative size of the box and dipenguin contribution shows that the latter is of the same order of magnitude as the box diagram – our estimate gives $r \sim 20 - 80\%$ depending on the choice of quark masses ($m_s = 0.1 \div 0.3 \text{ GeV}$).

The dipenguin diagram gives rise to a whole family of the long-distance contributions (e.g. Fig. 1c) that might be potentially important for $D^0 - \bar{D}^0$ mixing. In particular, it is plausible to assume that a gluon produced by a penguin (or any other operator) can form an intermediate bound state with two up-quarks, thus producing a hybrid resonance [6]. These one-particle intermediate state contributions to Δm_D might be of considerable importance because of the closure of mass

of hybrid resonance to the mass of D -meson. It is possible to show that a resonant long-distance contribution to Δm_D is given by ($D_S^L = 2^{-1/2} [D^0 \pm \bar{D}^0]$)

$$\Delta m_D = Re \frac{1}{2m_D} \sum_n \frac{|\langle D_L | H_w | n \rangle|^2 - |\langle D_S | H_w | n \rangle|^2}{m_D^2 - m_n^2 - \Gamma^2/4 + i\Gamma_n m_D}, \quad (10)$$

where we have neglected CP violation. The existence of the π_H or $\pi(1800) 0^{-+}$ meson was reported by several experimental groups [4] and predicted by the theoretical calculations in various quark models [5] suggesting a hybrid nature of this particle. One can estimate a contribution of $n = \pi_H$ to Eq.(10) provided that the mixing amplitude $g = \langle D_L | H_w | \pi_H \rangle$ is known. This amplitude can be estimated in various quark models, or phenomenologically, by using the available data on D decay rates. The idea is to seek for the common decay channels of D and π_H where π_H contribution is manifest and then estimate the mixing amplitude using the simple model for the $D - \pi_H$ mixing followed by π_H decay. It was noted in theoretical calculations [5] and hinted experimentally that the decay rates $\pi_H \rightarrow \pi f_0(980)$ and $\pi_H \rightarrow \pi f_0(1300)$ are large for the hybrid π_H . Thus one can put an upper bound on the mixing amplitude g by introducing a model for the resonant decay of D -meson via π_H :

$$\mathcal{M}(D \rightarrow \pi f_0(980)) = g (m_D^2 - m_{\pi_H}^2 + i\Gamma_{\pi_H} m_D)^{-1} A(\pi_H \rightarrow \pi f_0(980)). \quad (11)$$

Computing the mixing amplitude g using experimental data on the decay rate $\Gamma(D \rightarrow \pi f_0(980))$ and inserting it into Eq. (10), we estimate $\Delta m_D^{\pi_H} \leq 0.2 \times 10^{-16} \text{ GeV}$, which is five times larger than the short distance result.

In conclusion, we have shown that the short-distance dipenguin diagram contributes at the same order of magnitude as the box diagram to $D^0 - \bar{D}^0$ mixing amplitude. Long-distance dipenguin-like pieces presumably give larger contributions, but require further studies [2,6].

REFERENCES

1. A. Datta and D. Kumbhakar, Z. Phys. C **27**, 515 (1985); H.Y. Cheng, Phys. Rev. D**26**, 143 (1982)
2. J. Donoghue, E. Golowich, B. Holstein, and J. Trampetic, Phys. Rev. D **33**, 179 (1986); L. Wolfenstein, Phys. Lett. B **164**, 170 (1985); H. Georgi, *ibid.* **297**, 353 (1992); T. Ohl, G. Ricciardi, and E. Simmons, Nucl. Phys. **B403**, 605 (1993).
3. A.A. Petrov, Phys. Rev. **D56**, 1685 (1997).
4. D.V. Amelin, et.al. Phys. Lett. **356** (1995) 595.
5. F. Close, P. Page Nucl. Phys. **B443** (1995) 233; Phys. Rev. **D52** (1995) 1706.
6. A. Zaitsev, private communications; E. Golowich and A.A. Petrov, in preparation